

ORIGINAL RESEARCH ARTICLE

# Clinico-Aetiopathological Study of Erythroderma.

Chetana P.<sup>1</sup>, Krishnakanth M.<sup>2</sup>, Sudha R.<sup>3</sup>, Gayathri R.<sup>4</sup>, Murugan S.<sup>3</sup>,  
& Mahalakshmi V.<sup>5</sup>

<sup>1</sup>Junior Resident, <sup>2</sup>Associate Professor, <sup>3</sup>Professor, <sup>4</sup>Senior Resident, <sup>5</sup>Professor & HOD, Department of Dermatology, Venereology & Leprosy, Sri Ramachandra University, Porur, Chennai [TN], India.

## Abstract:

Erythroderma refers to any inflammatory skin disease affecting either sex or any age group resulting in erythema and exfoliation that affects more than 90% of the body surface. It may result from a pre-existing dermatoses or underlying lymphoma, drug eruption, hereditary causes i.e. ichthyosiform erythroderma. Hence it is mandatory to establish the cause in order to facilitate its management. Since it is a complex disorder, the patient should preferably be hospitalized for evaluation and treatment.

To study the clinical profile of patients with erythroderma and to identify the aetiological factors of erythroderma and its histopathological correlation and also its associated systemic complication. A total number of 57 clinically diagnosed cases of erythroderma attended the department during the period from 2012 to 2014 and were taken up for the clinico-aetiopathological evaluation.

Males out-numbered females with a ratio of 1.48 : 1. Predominant symptoms were erythroderma (96.49%), exfoliation (100%), itching (100%). Nail change were seen in 75.43% of patients. The most common underlying etiology observed was psoriasis (43.85%) followed by drug induced erythroderma (36.84%). The most common reaction pattern observed was psoriasiform pattern (59.6%). In 16 (33.33%) patients biopsy helped in reaching a diagnosis.

Erythroderma can be fatal, even when properly managed, primarily because of its metabolic complications. Hence it is mandatory to establish its aetiopathology in order to facilitate more practiced management apart from the routine basic management.

**Key Words:** Erythroderma, erythema, exfoliation, psoriasiform pattern.

## Introduction:

Erythroderma refers to any inflammatory skin disease affecting either sex or any age group resulting in erythema and exfoliation that affects more than 90% of the body surface. It may result from a pre-existing dermatoses or underlying lymphoma, drug eruption, hereditary causes eg: ichthyosiform erythroderma. Hence it is mandatory to establish the cause in order to facilitate its management.

The potential complications include secondary infection. Electrolyte imbalance, dehydration, temperature dysregulation, high output cardiac failure and death.

Since it is a complex disorder and is considered as one of the dermatologic emergency, the patient should preferably be hospitalized for evaluation and treatment.

## Materials and Methods:

To study the clinical profile of patients with erythroderma and to identify the aetiological factors of erythroderma and its histopathological correlation and also its associated systemic complication.

## Address for correspondence:

Dr. Krishnakanth M.

Associate Professor,  
Department of Dermatology,  
Sri Ramachandra University,  
Chennai [TN], India.

Email ID: krishnakanthmuralidhar@gmail.com

## How to cite this article:

Chetana P, Krishnakanth M, et al ; Clinico-Aetiopathological Study Of Erythroderma.; International Journal of current Medical and Applied sciences; 2015, 8(1), 01-05.

## Access this Article Online



Quick Response Code

Website:

[www.ijcmaas.com](http://www.ijcmaas.com)

Subject:


Medical Sciences

A total number of 57 clinically diagnosed cases of erythroderma attended in Department of Dermatology, Venereology & Leprosy during the period from December 2012 to September 2014 and were taken up for the clinio-aetiopathological evaluation. Patients of all ages and both the sexes who were willing to participate in the study were included. It was an Observational, Descriptive & Prospective hospital-based study. The materials included recording of participant demographics, documentation of clinical history, past medical and surgical history. The baseline investigations- Complete blood count, Renal and Liver function test, serum electrolytes, Urine microscopy, Chest X ray-P A View, Ultrasound-abdomen, ECG were done. Skin biopsy was done in 48 patients. Skin biopsy was not done in severely ill patients with altered

coagulation profile. Statistical Analysis was done using the Statistical package for the Social Sciences (SPSS V 17.1). Quantitative variable are expressed as mean ± standard Deviation. Qualitative variable are expressed as percentage values.

**Results:**

The most common underlying etiology of erythroderma found in the study was psoriasis (43.85%) followed by drug induced (36.84%). Other etiologies which were noted include air borne contact dermatitis (7.01%), pemphigus foliaceus (3.5%), non bullous ichthyosi form erythroderma (3.5%) and atopic dermatitis (1.75%). Etiology could not be identified in 2 patients (3.5%).




**Table 1 : Drug induced Erythroderma**

DRUG	% (n=57)
<b>Indigenous medications</b>	<b>14.03</b>
Anticonvulsants (Carbamazepine, Phenytoin)	12.28
Dapsone	5.26
ATT	1.75
Vincristine	1.75
Unrecordable drug	1.75


**Table 2: Common HPE Patterns**

Pattern	No.	Percentage
Psoriasiform	34	70.8
Lichenoid	7	14.5
Spongiotic	6	12.5
Vesiculo-bullous	1	2.1

Out of the 57 patients who were recruited in my study, skin biopsies were done in 48 patients. The most common reaction pattern observed was psoriasiform (70.8%). Other reaction patterns observed are lichenoid (14.5%), spongiotic (12.5%), and vesiculo- bullous (2.1%).



Nail changes however differed in present study .



Predominant symptoms –exfoliation of skin 100%, itching 100%, redness 96.49%, fever 80.70%, swelling of feet 56.14%, fever with chills 35.8% contact with allergen 7.1%, fluid filled lesions and atopy 5.26%

**Table 3: Complications seen in Erythroderma**

Complications	% (n=57)
Anemia	73.68
Leucocytosis	57.89
Eosinophilia	47.36
Elevated ESR	68.42
Altered RFT	15.78
Altered LFT	33.33
Hypoproteinemia	38.59
Hypoalbuminemia	40.35
Hepatomegaly (USG)	7.01

Haemoglobin was low in 42 cases (73.68%). Leucocytosis was noted in 57.89% of patients in this study. Eosinophilia was observed in 47.36% of patients. In this study, ESR was raised in 68.42% of patients. Hypoproteinemia and hypoalbuminemia occurred in 38.59% and 40.35% of cases in this study.

**Discussion:**

A total number of 57 clinically diagnosed cases of erythroderma attended in department during the period from December 2012 to September 2014 and

were taken up for the clinio-aetiopathological evaluation.

In this study, male outnumbered female in the ratio of 1.48 : 1. Earlier studies also demonstrated. Involvement of palms and soles were seen in 63.15% of patients in

this study while it was observed in 39.13% cases in a study conducted by Bharatiya PR et al [3] 70 and in 30% of cases in a study done by pal Haroon [4]. In this study, the mucosal involvement was recorded in 18 (31.57%) cases. However, in the study conducted by Pal S Haroon [4] the mucosal involvement was observed in 36.6% of cases and 1% in a study conducted by Akhyani et al [5].

In this study, psoriasis (43.85%) was the commonest cause of erythroderma. Similar findings were observed in studies conducted by Sehgal et al [5]. Pal S Haroon [4] Sudha et al [6].

Erythroderma due to underlying pemphigus foliaceus was seen in 3.5% of patients in this study. In studies conducted by Abrahams et al [7]. King Le et al [8] this dermatoses did not exceed 2%. In a study conducted by Pal S and Haroon, [4] the frequency of pemphigus foliaceus was 5.6 whereas 6.25% in a study conducted by Rym et al [9].

Erythroderma secondary to malignancy was not observed in this study. Similar findings were observed in a study conducted by Sehgal and Srivatsava [10], while Botella Estroda et al [11] reported 12.5% of erythroderma occurred due to cutaneous T-cell lymphoma and Kind Le et al [8] observed that 20% cases occurred due to lymphoreticulat neoplasms, especially cutaneous T-cell lymphoma.

In this study, 8.77% cases of erythroderma occurred secondary to various eczematous conditions, such as Air borne contact dermatitis (7.01%) and atopic dermatitis (1.75%). Erythroderma secondary to eczema was seen in 20% of cases in a study conducted by Chaundary et al [12].

In 3.5% of patients, the cause of erythroderma could not be found in this study. Similarly, in a study conducted by Rym et al [10] the cause could not be determined in 7.5% of patients and in 18% patients in another study conducted by Javeria et al [13] The cause could not be found in 14.6% of cases in a study done by Pal S and Haroon.

Among the 57 patients who were enrolled in this study, one patient (1.75%) was HIV- positive. The cause of erythroderma was due to unknown drug intake. There was no HIV – infected patients in a study conducted by Akhyani et al [5]. In a study conducted by Morar et al, drug reactions (40.6%) was the most common cause of erythroderma in HIV-positive patients and also concluded that in the young black patients erythroderma may be a marker for HIV infection [14].

During the study period, death occurred in two (3.5%) out of 57 patients due to fulminant hepatitis and cardiac failure. Similarly two patients died due to cardiac failure and septicaemia in a study conducted by Sudha et al [15]. In a study conducted by Rym et al, the

proportion of death was 3.75% [9] Erythroderma in adults: A report of 80 cases.

Nail changes however differed in our study from previous studies. In the study conducted by Sudha R [15] et al: pitting & onycholysis were the preponderance findings. Hulmani et al [1] demonstrated Ridging as their predominant findings. In our case elongation of nails was the predominant findings.

Systemic features such as Pedal edema was seen in 57.89% of cases in present study while in a study conducted by Hulmani M. et al [1], 70% of cases had pedal edema. Lymphadenopathy was seen in 23 cases (40.35%) in this study, whereas 55.5% of cases in pal and Haroon et al. [4 ] and 56.66% cases in Hulmani M. [1] et al. had lymphadenopathy.

Haemoglobin was low in 42 cases (73.68%). Leucocytosis was noted in 57.89% of patients in this study. Eosinophilia was observed in 47.36% of patients. In this study, ESR was raised in 68.42% of patients. Hypoproteinemia and hypoalbuminemia occurred in 38.59% and 40.35% of cases in this study. Hypoproteinemia with altered albumin to globulin was seen in 63.3% cases in a study conducted by Manjunath Hulmani et al.[1] and 12% in Sudha et al. [2]. Altered liver function tests was seen in 33.33% of cases in this study. In a study conducted by Bharatiya PR et al.[3], altered liver function tests was observed in 8.69% of cases. In this study, altered renal function tests were seen in 15.78% of cases while in a study conducted by Manjunath Hulmani et al [1], it was seen in 6.6% of patients.

### Conclusion:

Predominant symptoms were erythroderma (96.49%), exfoliation (100%), itching (100%). Nail change were seen in 75.43% of patients. The most common underlying etiology observed was psoriasis (43.85%) followed by drug induced erythroderma (36.84%). The most common reaction pattern observed was psoriasiform pattern (59.6%).

In 16 (33.33%) patients biopsy helped in reaching a diagnosis Erythroderma can be fatal, even when properly managed, primarily because of its metabolic complications. Hence it is mandatory to establish its aetiopathology in order to facilitate more practiced management apart from the routine basic management. The study will help in education patients with underlying disease (Ex. Psoriasis, atopic dermatitis) about possible triggers of erythroderma (irritants, abrupt discontinuation of certain therapies) in order to prevent erythroderma and to avoid medications & irritants that have previously caused erythroderma and also to create awareness among the medical

professionals regarding the medications that are potentially cross-reactive.

### References:

1. Hulmani M, NandaKishore B, Bhat M R, Sukumar D, Martis J, Kamath G, Srinath M K. Clinico-etiological study of 30 erythroderma cases from tertiary center in South India. *Indian Dermatol Online J* 2014;5:25-9.
2. Sudha R, Hussain S B, Bellraj E, Frederick M, Mahalaxmi V, Sobhana S, Anandan S. clinicopathological study of exfoliative dermatitis. *Indian J Dermatol Venereol Leprol* 2003;69:30-1.
3. Bharatiya PR, Joshi PB. Study of exfoliative dermatitis. *Indian J Dermatol Venereol Leprol* 1995;61:81-3
4. Pal S, Haroon TS. Erythroderma: A clinic-etiological study of 90 cases. *Int J Dermatol* 1998;37:104-7
5. Akhyani M, Ghodsi ZS, Toosi S, Dabbaghian H. Erythroderma: A clinical study of 97 cases. *BMC Dermatol* 2005;5:5.
6. Sudha R, Hussain S B, Bellraj E, Frederick M, Mahalaxmi V, Sobhana S, Anandan S. clinicopathological study of exfoliative dermatitis. *Indian J Dermatol Venereol Leprol* 2003;69:30-1
7. Abrahams I, McCarthy JT, Sanders SL. 101 cases of exfoliative dermatitis. *Arch Dermatol* 1963;8:96-101
8. King LE, Durfesne RG, Lovert G, et al. Erythroderma: review of 82 cases. *South Med J* 1986;79: 1210-1215
9. Rym BM, Mourad M, Bechir Z Dalenda E, Faika C, Iadh AM, et al. Erythroderma in adults: A report of 80 cases. *Int J Dermatol* 2005;44:731-5.
10. Sehgal VN, Srivastava G. Exfoliative dermatitis: A prospective study of 80 patients. *Dermatologica* 1986;173:278-84.
11. Botella-Estrada R, Sanmartin O, Oliver V, Febrer J, Aliaga A. Erythroderma: A clinicopathological study of 56 cases. *Arch Dermatol* 1994;130:1503-7
12. Chaudhary A, Gupte PD. Erythroderma: a study of incidence and aetiopathogenesis. *Indian J Dermatol Venereol Leprol* 1997;63: 38-9.
13. Javeria Hafeez, Zafar Iqbal Shaikh, Asher Ahmad Mashhood, Simeen-Ber-Rahman. Frequency of various etiological factors associated with erythroderma. *Journal of Pakistan Association of Dermatologist*. 2010; 20:70-74.
14. Morar N, Dlova N, Gupta AK, Naidoo DK, Aboobaker J, Ramdial PK. Erythroderma: A comparison between HIV positive and negative patients. *Int J Dermatol* 1999;38:895-900.
15. Sudha R, Hussain S B, Bellraj E, Frederick M, Mahalaxmi V, Sobhana S, Anandan S. Clinicopathological study of exfoliative dermatitis. *Indian J Dermatol Venereol Leprol* 2003; 69: 30-1.